

**Numerical Methods for Differential Equations FMNN10/NUMN12  
Exam 2015-08-20**

Exam duration 08:00 – 13:00. A minimum of 16 points out of 32 are required to pass. Your grade is determined by your score on this exam and your project scores from the fall of 2014, in accordance with the rules on the course home page.

No computers, pocket calculators, cell phones, browsing tablets or any other electronic devices, and no textbooks, lecture notes or written material, may be used during the exam.

1. (6p) For the initial value problem  $\dot{y} = f(y)$  the following 3-step method is proposed, in backward difference notation:

$$(\nabla + a\nabla^2 + b\nabla^3) y_n = hf(y_n).$$

- (a) Determine the parameters  $a$  and  $b$  such that the method's order of consistency is maximal. (3p)
- (b) Show that the resulting method is zero-stable. (3p)
2. (3p) A Runge–Kutta method has the stability function

$$R(z) = \frac{1 + 2z/3 + z^2/6}{1 - z/3},$$

where  $z = h\lambda$ .

- (a) Is the method implicit or explicit? Motivate your answer. (1p)
- (b) What is the minimum number of stages the method must have in order to have this stability function? (1p)
- (c) Determine whether the method is A-stable. (1p)
3. (5p) Consider the nonlinear two-point boundary value problem

$$\frac{d^2u}{dx^2} + au \frac{du}{dx} - u = 0$$

with one Neumann condition  $u'(0) = 0$  and one Dirichlet condition  $u(1) = 0$ , and where the parameter  $a$  is positive.

- (a) Introduce a suitable grid and discretize with a *standard second order method*. Give *all details about the grid*, such as the number of grid points and their location, as well as mesh width  $\Delta x$ , and formulate the discretization. Include the boundary conditions in the equation system. (3p)
- (b) The system of equations is nonlinear, and requires the use Newton's method. Construct the system's Jacobian matrix. (2p)

4. (2p) Consider the eigenvalue problem

$$u'' + u' = \lambda u$$

with Dirichlet conditions  $u(0) = u(1) = 0$ . Introduce a suitable grid and discretize with a *standard second order method*. Give *all details about the grid*, such as the number of grid points and their location, as well as mesh width  $\Delta x$ , and formulate the algebraic eigenvalue problem  $Au = \lambda u$  that results from the discretization. Specifically, construct the matrix  $A$ .

5. (5p) Let  $u(x)$  be a differentiable function on  $[0, 1]$  with periodic boundary conditions  $u(0) = u(1)$ , and let the inner product  $\langle \cdot, \cdot \rangle$  be defined by

$$\langle u, v \rangle = \int_0^1 uv \, dx.$$

- (a) Show that  $u'$  is orthogonal to  $u^p$ , for  $p \geq 1$ . (2p)
- (b) Consider the inviscid Burgers equation  $u_t - uu_x = 0$  with periodic boundary conditions and assume that the solution is continuously differentiable. Show that  $\|u(t, \cdot)\|_2$  remains constant for all  $t$ . (3p)
6. (5p) Classify the following PDEs for  $t \geq 0$  and  $x \in [0, 1]$ :

- (a)  $u_{yy} + u_{xx} = f(x, y)$
- (b)  $u_t - a \cdot u_x = 0$
- (c)  $u_t + \frac{1}{2}(u^2)_x = 0$
- (d)  $u_t = \frac{\partial}{\partial x} \left( p(x) \frac{\partial u}{\partial x} \right)$
- (e)  $u_t = \frac{1}{\text{Pe}} u_{xx} + u_x + f(t, x)$

For each equation, state whether the problem is *elliptic*, *parabolic* or *hyperbolic*. All parameters are supposed to be *positive*. In addition, give the "name" of each equation (e.g. "convection-diffusion equation" etc.).

7. (6p) Consider the convection–diffusion equation

$$u_t = u_{xx} + \gamma u_x$$

with homogeneous boundary conditions, and initial condition  $u(0, x) = g(x)$ . The differential operator

$$\mathcal{L} = \frac{\partial^2}{\partial x^2} + \gamma \frac{\partial}{\partial x}$$

then has negative real eigenvalues,

$$\lambda_k = -(k\pi)^2 - \frac{\gamma^2}{4}.$$

- (a) Introduce a grid and use a symmetric second order discretization in space. Write down your discretization in matrix-vector form. (2p)
- (b) Given that the eigenvalues of an non-symmetric tridiagonal  $N \times N$  Toeplitz matrix  $A = \text{tridiag}(b \ a \ c)$  are

$$\lambda_k = a + 2\sqrt{bc} \cos \frac{k\pi}{N+1}; \quad k = 1 : N,$$

what is the largest value of  $\Delta x$  you can choose so that the discrete problem also has negative real eigenvalues? (2p)

- (c) Combine your method of lines discretization with the explicit Euler method for time stepping. What is the CFL condition on  $\Delta t$  in order to guarantee stability? (2p)

GOOD LUCK!  
G.S.